

Patient: _____ Date: _____ Time: _____ AM/PM

The Timed Up and Go (TUG) Test

Purpose: To assess mobility

Equipment: A stopwatch

Directions: Patients wear their regular footwear and can use a walking aid if needed. Begin by having the patient sit back in a standard arm chair and identify a line 3 meters or 10 feet away on the floor.

Instructions to the patient:

When I say **"Go,"** I want you to:

1. Stand up from the chair
2. Walk to the line on the floor at your normal pace
3. Turn
4. Walk back to the chair at your normal pace
5. Sit down again

On the word **"Go"** begin timing.

Stop timing after patient has sat back down and record.

Time: _____ **seconds**

An older adult who takes ≥ 12 seconds to complete the TUG is at high risk for falling.

Observe the patient's postural stability, gait, stride length, and sway.

Circle all that apply: Slow tentative pace Loss of balance
Short strides Little or no arm swing Steadying self on walls
Shuffling En bloc turning Not using assistive device properly

Notes:

For relevant articles, go to: www.cdc.gov/injury/STEADI



Centers for Disease
Control and Prevention
National Center for Injury
Prevention and Control

STEADI Stopping Elderly
Accidents, Deaths & Injuries

Patient: _____ Date: _____ Time: _____ AM/PM

The 30-Second Chair Stand Test


Purpose: To test leg strength and endurance

Equipment:

- A chair with a straight back without arm rests (seat 17" high)
- A stopwatch

Instructions to the patient:

1. Sit in the middle of the chair.
2. Place your hands on the opposite shoulder crossed at the wrists.
3. Keep your feet flat on the floor.
4. Keep your back straight and keep your arms against your chest.
5. On **"Go,"** rise to a full standing position and then sit back down again.
6. Repeat this for 30 seconds.



On **"Go,"** begin timing.

If the patient must use his/her arms to stand, stop the test. Record "0" for the number and score.

Count the number of times the patient comes to a full standing position in 30 seconds.

If the patient is over halfway to a standing position when 30 seconds have elapsed, count it as a stand.

Record the number of times the patient stands in 30 seconds.

Number: _____ **Score** _____ **See next page.**

A below average score indicates a high risk for falls.

Notes:

For relevant articles, go to: www.cdc.gov/injury/STEADI



Centers for Disease
Control and Prevention
National Center for Injury
Prevention and Control

STEADI Stopping Elderly
Accidents, Deaths & Injuries

Chair Stand—Below Average Scores

Age	Men	Women
60-64	< 14	< 12
65-69	< 12	< 11
70-74	< 12	< 10
75-79	< 11	< 10
80-84	< 10	< 9
85-89	< 8	< 8
90-94	< 7	< 4